

**Solvipurity**

ANALYTICAL LABORATORY · REYKJAVÍK, IS

SVP-2026-00283ISSUED 2026-03-16 · ACCREDITATION AL-1142
ISO/IEC 17025 · GMP · GLP


CERTIFICATE OF ANALYSIS

**AUTHENTIC**

B7-33 6mg

Björn Healthcare ehf. · Sterile lyophilizate, 6 mg per 3 ml clear glass vial, rubber s
topper + aluminium flip-off (white cake)

REPRESENTATIVE CHROMATOGRAM · HPLC-UV 205 NM



BATCH NO.

BJRN-209Q96T

ANALYTICAL METHODS

RP-HPLC-UV 220 nm · LC-ESI-MS · AAA
(amino acid analysis) · Ion chromatography
(counter-ion) · Karl Fischer 2.5.32 · GC-MS
(headspace) · ICP-MS · Kinetic chromogenic
LAL 2.6.14 · Ph. Eur. 2.6.1 sterility · Ph. Eur. 2.6.12
microbial limits

MANUFACTURED

2026-02-12

EXPIRY

2027-11-12

RECEIVED

2026-03-16

RELEASE

2026-03-16

DECLARED COMPOSITION

B7-33 6mg — synthetic peptide; sequence: relaxin-2 B-chain single-chain analog (33 aa); CAS n/a;
theoretical MW 3251.70 Da

Analytical results

19 TESTS · ALL METHODS VALIDATED

| SUBSTANCE / PARAMETER | RESULT | LOQ | LIMIT | METHOD |
|---|---|-----|---|-------------------------|
| ● Appearance — sterile lyophilized cake, white to off-white | Conforms | — | homogeneous white cake, no particulates | Visual |
| ● Solubility (water for injection, 2 mg/ml, 25 °C) | Complete within 60 s — clear colourless solution | — | Clear, no visible particles | Visual (Ph. Eur. 2.2.1) |

| Identification — HPLC retention time | Matches reference | — | ±2.0 % of ref | RP-HPLC-UV 220 nm SVP-2026-00283 + |
|--|-----------------------------------|-------------|-----------------------------------|---|
| ● Identification — sequence / mass match | Confirmed CAS n/a | — | Match theoretical within ±1 Da | LC-ESI-MS |
| ● Molecular weight (measured) | 3251.54 Da Δ = -0.16 Da | 0.5 Da | 3251.70 Da ± 1.0 Da (theoretical) | ESI-MS |
| ● Chromatographic purity (main peak) | 99.35 % | 0.05 % | ≥ 98.0 % | RP-HPLC-UV 220 nm |
| ● Any single impurity (max) | 0.23 % | 0.05 % | ≤ 1.00 % | RP-HPLC-UV 220 nm |
| ● Peptide content (amino acid analysis) | 86.8 % w/w | 0.5 % | ≥ 80.0 % w/w | AAA (6 N HCl, 110 °C, 24 h) |
| ● Trifluoroacetate (TFA counter-ion) | 0.38 % w/w | 0.05 % | ≤ 1.00 % w/w | IC (ion chromatography) |
| ● Water content (Karl Fischer) | 3.55 % w/w | 0.1 % | ≤ 5.0 % w/w | Ph. Eur. 2.5.32 |
| ● Residual acetonitrile | 166 ppm | 10 ppm | ≤ 410 ppm (ICH Q3C Class 2) | GC-MS (headspace) |
| ● Residual DMF | 55 ppm | 10 ppm | ≤ 880 ppm (ICH Q3C Class 2) | GC-MS (headspace) |
| ● Lead (Pb) | 0.032 ppm | 0.02 ppm | ≤ 0.5 ppm (ICH Q3D parenteral) | ICP-MS |
| ● Arsenic + Cadmium + Mercury (total) | 0.172 ppm | 0.02 ppm | ≤ 1.5 ppm (ICH Q3D parenteral) | ICP-MS |
| ● Bacterial endotoxins (LAL) | 2.93 EU/mg | 0.125 EU/mg | < 10.0 EU/mg | Kinetic chromogenic LAL (Ph. Eur. 2.6.14) |
| ● TAMC (aerobic bacteria, pre-lyophilization bulk) | 8 CFU/g | 1 CFU/g | ≤ 10 ² CFU/g | Ph. Eur. 2.6.12 |
| ● TYMC (yeast / molds, pre-lyophilization bulk) | 3 CFU/g | 1 CFU/g | ≤ 10 ¹ CFU/g | Ph. Eur. 2.6.12 |
| ● Sterility (final lyophilized vial) | Complies — no growth | — | No growth, 14 d incubation | Ph. Eur. 2.6.1 (direct inoculation) |
| ● Container closure integrity | Pass | — | No dye uptake | Dye ingress (0.05 % methylene blue, 2 h vacuum) |

