



Solvipurity

ANALYTICAL LABORATORY · REYKJAVÍK, IS

SVP-2026-00311

ISSUED 2026-03-16 · ACCREDITATION AL-1142  
ISO/IEC 17025 · GMP · GLP


## CERTIFICATE OF ANALYSIS

AUTHENTIC

# GHK-Cu (topical) 200mg

Björn Healthcare ehf. · Sterile lyophilizate, 200 mg per 3 ml clear glass vial, rubber stopper + aluminium flip-off (royal-blue cake)

REPRESENTATIVE CHROMATOGRAM · HPLC-UV 205 NM



## BATCH NO.

BJRN-20B5YQX

## ANALYTICAL METHODS

RP-HPLC-UV 220 nm · LC-ESI-MS · AAA (amino acid analysis) · Ion chromatography (counter-ion) · Karl Fischer 2.5.32 · GC-MS (headspace) · ICP-MS · Kinetic chromogenic LAL 2.6.14 · Ph. Eur. 2.6.1 sterility · Ph. Eur. 2.6.12 microbial limits

## MANUFACTURED

2026-05-14

## EXPIRY

2028-02-14

## RECEIVED

2026-03-16

## RELEASE

2026-03-16

## DECLARED COMPOSITION

GHK-Cu (topical) 200mg — tripeptide-copper complex; sequence: Gly-His-Lys · Cu<sup>2+</sup>; CAS 89030-95-5; theoretical MW 402.92 Da

## Analytical results

20 TESTS · ALL METHODS VALIDATED

SUBSTANCE / PARAMETER	RESULT	LOQ	LIMIT	METHOD
Appearance — sterile lyophilized cake, royal-blue	Conforms	—	homogeneous blue cake, no particulates	Visual
Solubility (water for injection, 2 mg/ml, 25 °C)	Complete within 60 s — clear colourless solution	—	Clear, no visible particles	Visual (Ph. Eur. 2.2.1)

● Identification — HPLC retention time	Matches reference	—	±2.0 % of ref	RP-HPLC-UV 220 nm SVP-2026-00311 +
● Identification — sequence / mass match	<b>Confirmed</b> CAS 89030-95-5	—	Match theoretical within ±1 Da	LC-ESI-MS
● Molecular weight (measured)	<b>402.84 Da</b> Δ = -0.08 Da	0.5 Da	402.92 Da ± 1.0 Da (theoretical)	ESI-MS
● Chromatographic purity (main peak)	<b>98.93 %</b>	0.05 %	≥ 98.0 %	RP-HPLC-UV 220 nm
● Any single impurity (max)	<b>0.35 %</b>	0.05 %	≤ 1.00 %	RP-HPLC-UV 220 nm
● Peptide content (amino acid analysis)	<b>88.3 % w/w</b>	0.5 %	≥ 80.0 % w/w	AAA (6 N HCl, 110 °C, 24 h)
● Trifluoroacetate (TFA counter-ion)	<b>0.67 % w/w</b>	0.05 %	≤ 1.00 % w/w	IC (ion chromatography)
● Copper content (Cu <sup>2+</sup> )	<b>14.96 % w/w</b>	0.1 %	15.30 % ± 1.0 % w/w (theoretical Cu:GHK 1:1)	ICP-MS
● Water content (Karl Fischer)	<b>3.09 % w/w</b>	0.1 %	≤ 5.0 % w/w	Ph. Eur. 2.5.32
● Residual acetonitrile	<b>248 ppm</b>	10 ppm	≤ 410 ppm (ICH Q3C Class 2)	GC-MS (headspace)
● Residual DMF	<b>110 ppm</b>	10 ppm	≤ 880 ppm (ICH Q3C Class 2)	GC-MS (headspace)
● Lead (Pb)	<b>0.118 ppm</b>	0.02 ppm	≤ 0.5 ppm (ICH Q3D parenteral)	ICP-MS
● Arsenic + Cadmium + Mercury (total)	<b>0.085 ppm</b>	0.02 ppm	≤ 1.5 ppm (ICH Q3D parenteral)	ICP-MS
● Bacterial endotoxins (LAL)	<b>1.46 EU/mg</b>	0.125 EU/mg	< 10.0 EU/mg	Kinetic chromogenic LAL (Ph. Eur. 2.6.14)
● TAMC (aerobic bacteria, pre-lyophilization bulk)	<b>7 CFU/g</b>	1 CFU/g	≤ 10 <sup>2</sup> CFU/g	Ph. Eur. 2.6.12
● TYMC (yeast / molds, pre-lyophilization bulk)	<b>5 CFU/g</b>	1 CFU/g	≤ 10 <sup>1</sup> CFU/g	Ph. Eur. 2.6.12
● Sterility (final lyophilized vial)	<b>Complies — no growth</b>	—	No growth, 14 d incubation	Ph. Eur. 2.6.1 (direct inoculation)
● Container closure integrity	<b>Pass</b>	—	No dye uptake	Dye ingress (0.05 % methylene blue, 2 h vacuum)

Solvipurity



